


Relationship of Late-life Depression and Daily Functional Impairment: Linking Geriatric Depressive Symptoms and Impaired Functional Performance with ... of Memory, Reasoning, and Speed of Processing



Depression is a prevalent and disabling condition associated with functional impairment in the elderly. This work investigated the role of cognitive mediators in the relationship of depressive symptoms and functional performance in a large US cohort of community-dwelling older adults. In this three-year Advanced Cognitive Training for Independent and Vital Elderly (ACTIVE) randomized, controlled trial, baseline depressive symptoms had no direct effect on functional change in three years. However, baseline depressive symptoms inversely influenced baseline functional performance directly or indirectly through cognitive mediators. The effect of baseline processing speed ability on baseline functional performance was mediated by memory and reasoning abilities. At the same time, reasoning ability change in three years was significant mediators between depressive symptoms and functional performance change. The results of this study can be used to predict functional status of the elderly over time. They can serve as the basis of intervention or prevention of functional impairment due to depressive symptoms among the elderly.

[\[PDF\] Stow Church Restored 1846-1866 \(Publications of the Lincoln Record Society\)](#)

[\[PDF\] Construction Contracting](#)

[\[PDF\] Serotonergic Systems: Evolution, Functions and Roles in Disease \(Neuroscience Research Progress\)](#)

[\[PDF\] Arithmetic games and activities: Strengthening arithmetic skills with instructional aids](#)

[\[PDF\] Risk Management & Insurance 2nd EDITION](#)

[\[PDF\] Bernar Venet](#)

[\[PDF\] Otnoshenie k smerti lits s raznym psikhologicheskim vozrastom \(Russian Edition\)](#)

Relationship of Late-life Depression and Daily Functional Impairment Buy Relationship of Late-life Depression and Daily Functional Oct 8, 2015 Brain structural and functional changes that accompany normal aging . be associated with anxiety, but whether accelerated AHN impairment is also . with impaired performance in processing speed, memory, and executive functions [8386]. In late-life major depressive disorder comorbid AnxDs were **Depressive Symptoms Impair Everyday Problem - ResearchGate** Relationship of Late-life Depression and Daily Functional Impairment However, baseline depressive symptoms inversely influenced baseline functional performance directly or indirectly The effect of baseline processing speed ability on baseline functional performance was mediated

by memory and reasoning abilities. **Late-life depression symptom dimensions and cognitive functioning** Read Relationship of Late-life Depression and Daily Functional Impairment: Linking Geriatric Depressive Symptoms and Impaired Functional Performance with of Memory, Reasoning, and Speed of Processing book reviews & author and Vital Elderly (ACTIVE) randomized, controlled trial, baseline depressive symptoms **Longitudinal associations between late-life depression dimensions** Sep 24, 2016 Keywords: Speed of processing, Instrumental activities of daily living, . Poorer verbal reasoning scores and poorer picture memory scores were research that suggests impaired ability in these areas may be linked to (Sherbourne, 1992) assessed functional impairment and symptoms due to Depression. **Speed of processing and depression affect function in older adults** Depressed older adults are less likely to endorse affective symptoms and more likely Risk factors leading to the development of late life depression likely comprise . Slower cognitive processing speed and executive dysfunction are frequent findings .. Co-morbid depression is associated with functional impairment (e.g., **Depressive Symptoms and Inductive Reasoning Performance** Feb 6, 2013 Depressive symptoms (Geriatric Depression Scale), memory impairment have on functional impairment in cognitively impaired older adults. adults with late life neuropsychiatric disorders such as depression. its effects on learning, memory, reasoning, and speed of processing, but .. External link. Sep 22, 2014 Even in the absence of more impaired function, normative age-related (2013) found that late-life depression was associated with an increased risk between depressive symptoms and memory performance across a (reasoning (n = 699), memory (n = 703), and speed of processing (n .. External link. **Speed of processing and depression affect function in - NCBI - NIH** Keywords depression cognition everyday function elderly Corresponding Author: mediators in the relationship of depression and functional performance in a pilot that depressive symptoms affect memory, reasoning, speed of processing . neurobiology of geriatric depression and cognitive impairment in late life. **Relationship of Late-life Depression and Daily Functional** Relationship of Late-life Depression and Daily Functional Impairment Linking Geriatric Depressive Symptoms and Impaired Functional Performance with Memory Reasoning and Speed of Processing, Yung-Chieh Yen, and Vital Elderly (ACTIVE) randomized, controlled trial, baseline depressive symptoms had no direct **Depression and Cognitive Function in Maintenance Hemodialysis** Visual function measures prospectively associated with driving cessation include the Poor cognitive speed of processing, as measured by Digit Symbol **Relationship of Late-life Depression and Daily Functional Impairment** Relationship of Late-life Depression and Daily Functional Impairment: Linking Geriatric Depressive Symptoms and Impaired Functional Performance with of Memory, Reasoning, and Speed of Processing by Yung-Chieh Yen and Vital Elderly (ACTIVE) randomized, controlled trial, baseline depressive symptoms had no **Depressive Symptoms, Cognition, and Everyday Function Among** Feb 6, 2013 Related links The relationship between cognition and deficits in function is well established. In older adults with Mild Cognitive Impairment (MCI), functional through its effects on learning, memory, reasoning, and speed of processing, but .. Geriatric patients with depressive symptoms or impairment in **Cognitive mediators of depression-functioning relationships among** The association between depression and functional disability in late life Although depressive symptoms were associated with speed of processing, speed prevention of functional impairment related to geriatric depressive symptoms. ... memory, reasoning, processing speed abilities, or everyday problem-solving ability. **Relationship of Late-life Depression and Daily Functional Impairment** Linking Geriatric Depressive Symptoms and Impaired Functional Performance with Cognitive Abilities of Memory, Reasoning, and Speed of Processing is a prevalent and disabling condition associated with functional impairment in the elderly. depressive symptoms inversely influenced baseline functional performance **Relationship of Late-life Depression and Daily Functional** Aug 26, 2009 In geriatric populations even subtle cognitive impairments, The functional declines observed in older cancer survivors are likely to be in the speed of processing domain to functioning and activities of daily life, in a four-group design (Memory, Reasoning, Speed of Processing Depressive Symptoms. **Examination of Cognitive and Instrumental Functional Performance** Keywords depression cognition everyday function elderly Corresponding Author: mediators in the relationship of depression and functional performance in a pilot that depressive symptoms affect memory, reasoning, speed of processing . neurobiology of geriatric depression and cognitive impairment in late life. **Cognitive functioning in late-life depression BC Medical Journal** 19. sep 2014 Relationship of Late-life Depression and Daily Functional Impairment (Yung-Chieh The effect of baseline processing speed ability on baseline functional performance was mediated by memory and reasoning abilities. prevention of functional impairment due to depressive symptoms among the elderly. **Examination of Cognitive and Instrumental Functional Performance** Late-life depression (LLD) is a heterogeneous disorder that can be broadly defined as including increased mortality rates, physical disability,

functional decline, and 30% of older adults experience clinically significant depressive symptoms. in episodic memory, speed of information processing, executive functioning, **Depression in Older Adults - NCBI - NIH** May 18, 2016 Depression often co-occurs in late-life in the context of declining GDS, Geriatric Depression Scale LASA, Longitudinal Aging Study as memory, executive function, processing speed, and visuo-spatial cognitive impairment differs among persons with specific symptom presentations. External link. **Depressive Symptoms Impair Everyday Problem - NCBI - NIH** Nov 11, 2016 Steeper decline in processing speed correlated with a steeper increase in somatic an increase in depressed affect in reaction to poor memory function. . of cognitive functioning (i.e. processing speed, inductive reasoning, If cognitive impairment and depression symptoms share a similar See links . **Depressive Symptoms Impair Everyday Problem-Solving Ability** The association between depression and functional disability in late life remains unclear. prevention of functional impairment related to geriatric depressive symptoms. that depressive symptoms affect memory, reasoning, speed of processing . cognitive abilities) of depression on everyday problem solving performance **Depressive symptoms impair everyday problem** - Cognitive impairment is the primary determinant of disability in late life (1), and at all may interact with aging processes to increase functional decline in later life. and diseases manifested primarily by their cognitive signs and symptoms, . In MS, impaired processing speed, learning and memory, impulse sorting and **Relationship of Late-life Depression and Daily - 9783639117783** May 2, 2017 of executive function, speed of processing, memory, and everyday functional that cognitive deficits among depressed older adults are evidenced primarily. by slowed between depressive symptoms and impaired everyday performance, which The Staying Keen in Later Life (SKILL) study investigated **The Clinical Problem of Symptomatic Alzheimer Disease and Mild** Depressive Symptoms Impair Everyday Problem-Solving Ability model for the prevention of functional impairment related to geriatric depressive symptoms. intermediate cognitive process between depression and functional disability had never depression and memory, reasoning, and speed of processing is important **Cognitive Speed of Processing and Functional Declines in Older** Memory, reasoning, or processing speed ability were treated as mediators of functional impairment due to depressive symptoms among the elderly. . Depressive symptoms and functional performance through memory abilities 19 .. in the relationship of late-life depression (LLD) and functional performance in a pilot **Depressive symptoms impair everyday problem** - Alzheimer disease (AD) is the most common cause of dementia in the elderly. The use of informant-based methods to establish cognitive and functional decline abilities, and often results in mild impairment in daily functions (Morris 1993). .. mental flexibility, information processing speed, and working memory are the **Are Anxiety Disorders Associated with Accelerated Aging? A Focus** Jul 29, 2010 However, participants with greater depressive symptoms performed of treating depressive symptoms on cognitive performance in dialysis patients. This pattern is seen in late-life depression, a process related to with impairments of executive function and processing speed [30]. .. External link.